

# Interpretation of Test Results for HCV Infection and Further Actions

TEST OUTCOME	INTERPRETATION	FURTHER ACTIONS
HCV antibody nonreactive	No HCV antibody detected	Sample can be reported as nonreactive for HCV antibody. No further action required.  If recent exposure in person tested is suspected, test for HCV RNA.*
HCV antibody reactive	Presumptive HCV infection	A repeatedly reactive result is consistent with current HCV infection, or past HCV infection that has resolved, or biologic false positivity for HCV antibody. Test for HCV RNA to identify current infection.
HCV antibody reactive, HCV RNA detected	Current HCV infection	Provide person tested with appropriate counseling and link person tested to care and treatment.†
HCV antibody reactive, HCV RNA not detected	No current HCV infection	No further action required in most cases.  If distinction between true positivity and biologic false positivity for HCV antibody is desired, and if sample is repeatedly reactive in the initial test, test with another HCV antibody assay.  In certain situations,‡ follow up with HCV RNA testing and appropriate counseling.

\* If HCV RNA testing is not feasible and person tested is not immunocompromised, do follow-up testing for HCV antibody to demonstrate seroconversion. If the person tested is immunocompromised, consider testing for HCV RNA.

† It is recommended before initiating antiviral therapy to retest for HCV RNA in a subsequent blood sample to confirm HCV RNA positivity.

‡ If the person tested is suspected of having HCV exposure within the past 6 months, or has clinical evidence of HCV disease, or if there is concern regarding the handling or storage of the test specimen.