

children require more immediate attention for elevated BP. The difference between the 95th and 99th percentiles is only 7–10 mmHg and is not large enough, particularly in view of the variability in BP measurements, to adequately distinguish mild hypertension—where limited evaluation is most appropriate—from more severe hypertension where more immediate and extensive intervention is indicated.

Therefore, Stage 1 hypertension is the designation for BP levels that range from the 95th percentile to 5 mmHg above the 99th percentile. Stage 2 hypertension is the designation for BP levels that are higher than 5 mmHg above the 99th percentile. Once confirmed on repeated measures, Stage 1 hypertension allows time for evaluation before initiating treatment unless the patient is symptomatic. Patients with Stage 2 hypertension may need more prompt evaluation and pharmacologic therapy. Symptomatic patients with Stage 2 hypertension require immediate treatment and consultation with experts in pediatric hypertension. These categories are parallel to the staging of hypertension in adults, as noted in JNC 7.<sup>2</sup>

### USING THE BLOOD PRESSURE TABLES

1. Use the standard height charts to determine the height percentile.
2. Measure and record the child's SBP and DBP.
3. Use the correct gender table for SBP and DBP.
4. Find the child's age on the left side of the table. Follow the age row horizontally across the table to the intersection of the line for the height percentile (vertical column).
5. There, find the 50th, 90th, 95th, and 99th percentiles for SBP in the left columns and for DBP in the right columns.
  - BP less than the 90th percentile is normal.
  - BP between the 90th and 95th percentile is prehypertension. In adolescents, BP equal to or exceeding 120/80 mmHg is prehypertension, even if this figure is less than the 90th percentile.
  - BP greater than the 95th percentile may be hypertension.
6. If the BP is greater than the 90th percentile, the BP should be repeated twice at the same office visit, and an average SBP and DBP should be used.
7. If the BP is greater than the 95th percentile, BP should be staged. If Stage 1 (95th percentile to the 99th percentile plus 5 mmHg), BP measurements should be repeated on two more occasions. If hypertension is confirmed, evaluation should proceed as described in table 7. If BP is Stage 2 (>99th percentile plus 5 mmHg), prompt referral should be made for evaluation and therapy. If the patient is symptomatic, immediate referral and treatment are indicated. Those patients with a compelling indication, as noted in table 6, would be treated as the next higher category of hypertension.

TABLE 3

## Blood Pressure Levels for Boys by Age and Height Percentile\*

Age (Year)	BP Percentile ↓	Systolic BP (mmHg)							Diastolic BP (mmHg)						
		← Percentile of Height →							← Percentile of Height →						
		5th	10th	25th	50th	75th	90th	95th	5th	10th	25th	50th	75th	90th	95th
1	50th	80	81	83	85	87	88	89	34	35	36	37	38	39	39
	90th	94	95	97	99	100	102	103	49	50	51	52	53	53	54
	95th	98	99	101	103	104	106	106	54	54	55	56	57	58	58
	99th	105	106	108	110	112	113	114	61	62	63	64	65	66	66
2	50th	84	85	87	88	90	92	92	39	40	41	42	43	44	44
	90th	97	99	100	102	104	105	106	54	55	56	57	58	58	59
	95th	101	102	104	106	108	109	110	59	59	60	61	62	63	63
	99th	109	110	111	113	115	117	117	66	67	68	69	70	71	71
3	50th	86	87	89	91	93	94	95	44	44	45	46	47	48	48
	90th	100	101	103	105	107	108	109	59	59	60	61	62	63	63
	95th	104	105	107	109	110	112	113	63	63	64	65	66	67	67
	99th	111	112	114	116	118	119	120	71	71	72	73	74	75	75
4	50th	88	89	91	93	95	96	97	47	48	49	50	51	51	52
	90th	102	103	105	107	109	110	111	62	63	64	65	66	66	67
	95th	106	107	109	111	112	114	115	66	67	68	69	70	71	71
	99th	113	114	116	118	120	121	122	74	75	76	77	78	78	79
5	50th	90	91	93	95	96	98	98	50	51	52	53	54	55	55
	90th	104	105	106	108	110	111	112	65	66	67	68	69	69	70
	95th	108	109	110	112	114	115	116	69	70	71	72	73	74	74
	99th	115	116	118	120	121	123	123	77	78	79	80	81	81	82
6	50th	91	92	94	96	98	99	100	53	53	54	55	56	57	57
	90th	105	106	108	110	111	113	113	68	68	69	70	71	72	72
	95th	109	110	112	114	115	117	117	72	72	73	74	75	76	76
	99th	116	117	119	121	123	124	125	80	80	81	82	83	84	84
7	50th	92	94	95	97	99	100	101	55	55	56	57	58	59	59
	90th	106	107	109	111	113	114	115	70	70	71	72	73	74	74
	95th	110	111	113	115	117	118	119	74	74	75	76	77	78	78
	99th	117	118	120	122	124	125	126	82	82	83	84	85	86	86
8	50th	94	95	97	99	100	102	102	56	57	58	59	60	60	61
	90th	107	109	110	112	114	115	116	71	72	72	73	74	75	76
	95th	111	112	114	116	118	119	120	75	76	77	78	79	79	80
	99th	119	120	122	123	125	127	127	83	84	85	86	87	87	88
9	50th	95	96	98	100	102	103	104	57	58	59	60	61	61	62
	90th	109	110	112	114	115	117	118	72	73	74	75	76	76	77
	95th	113	114	116	118	119	121	121	76	77	78	79	80	81	81
	99th	120	121	123	125	127	128	129	84	85	86	87	88	88	89
10	50th	97	98	100	102	103	105	106	58	59	60	61	61	62	63
	90th	111	112	114	115	117	119	119	73	73	74	75	76	77	78
	95th	115	116	117	119	121	122	123	77	78	79	80	81	81	82
	99th	122	123	125	127	128	130	130	85	86	86	88	88	89	90

TABLE 4

## Blood Pressure Levels for Girls by Age and Height Percentile\*

Age (Year)	BP Percentile ↓	Systolic BP (mmHg)							Diastolic BP (mmHg)						
		← Percentile of Height →							← Percentile of Height →						
		5th	10th	25th	50th	75th	90th	95th	5th	10th	25th	50th	75th	90th	95th
1	50th	83	84	85	86	88	89	90	38	39	39	40	41	41	42
	90th	97	97	98	100	101	102	103	52	53	53	54	55	55	56
	95th	100	101	102	104	105	106	107	56	57	57	58	59	59	60
	99th	108	108	109	111	112	113	114	64	64	65	65	66	67	67
2	50th	85	85	87	88	89	91	91	43	44	44	45	46	46	47
	90th	98	99	100	101	103	104	105	57	58	58	59	60	61	61
	95th	102	103	104	105	107	108	109	61	62	62	63	64	65	65
	99th	109	110	111	112	114	115	116	69	69	70	70	71	72	72
3	50th	86	87	88	89	91	92	93	47	48	48	49	50	50	51
	90th	100	100	102	103	104	106	106	61	62	62	63	64	64	65
	95th	104	104	105	107	108	109	110	65	66	66	67	68	68	69
	99th	111	111	113	114	115	116	117	73	73	74	74	75	76	76
4	50th	88	88	90	91	92	94	94	50	50	51	52	52	53	54
	90th	101	102	103	104	106	107	108	64	64	65	66	67	67	68
	95th	105	106	107	108	110	111	112	68	68	69	70	71	71	72
	99th	112	113	114	115	117	118	119	76	76	76	77	78	79	79
5	50th	89	90	91	93	94	95	96	52	53	53	54	55	55	56
	90th	103	103	105	106	107	109	109	66	67	67	68	69	69	70
	95th	107	107	108	110	111	112	113	70	71	71	72	73	73	74
	99th	114	114	116	117	118	120	120	78	78	79	79	80	81	81
6	50th	91	92	93	94	96	97	98	54	54	55	56	56	57	58
	90th	104	105	106	108	109	110	111	68	68	69	70	70	71	72
	95th	108	109	110	111	113	114	115	72	72	73	74	74	75	76
	99th	115	116	117	119	120	121	122	80	80	80	81	82	83	83
7	50th	93	93	95	96	97	99	99	55	56	56	57	58	58	59
	90th	106	107	108	109	111	112	113	69	70	70	71	72	72	73
	95th	110	111	112	113	115	116	116	73	74	74	75	76	76	77
	99th	117	118	119	120	122	123	124	81	81	82	82	83	84	84
8	50th	95	95	96	98	99	100	101	57	57	57	58	59	60	60
	90th	108	109	110	111	113	114	114	71	71	71	72	73	74	74
	95th	112	112	114	115	116	118	118	75	75	75	76	77	78	78
	99th	119	120	121	122	123	125	125	82	82	83	83	84	85	86
9	50th	96	97	98	100	101	102	103	58	58	58	59	60	61	61
	90th	110	110	112	113	114	116	116	72	72	72	73	74	75	75
	95th	114	114	115	117	118	119	120	76	76	76	77	78	79	79
	99th	121	121	123	124	125	127	127	83	83	84	84	85	86	87
10	50th	98	99	100	102	103	104	105	59	59	59	60	61	62	62
	90th	112	112	114	115	116	118	118	73	73	73	74	75	76	76
	95th	116	116	117	119	120	121	122	77	77	77	78	79	80	80
	99th	123	123	125	126	127	129	129	84	84	85	86	86	87	88

TABLE 5

**Classification of Hypertension in Children and Adolescents,  
With Measurement Frequency and Therapy Recommendations**

	SBP or DBP Percentile*	Frequency of BP Measurement	Therapeutic Lifestyle Changes	Pharmacologic Therapy
Normal	<90th	Recheck at next scheduled physical examination.	Encourage healthy diet, sleep, and physical activity.	—
Prehypertension	90th to <95th or if BP exceeds 120/80 mmHg even if below 90th percentile up to <95th percentile†	Recheck in 6 months.	Weight-management counseling if overweight, introduce physical activity and diet management.‡	None unless compelling indications such as CKD, diabetes mellitus, heart failure, or LVH exist
Stage 1 hypertension	95th percentile to the 99th percentile plus 5 mmHg	Recheck in 1–2 weeks or sooner if the patient is symptomatic; if persistently elevated on two additional occasions, evaluate or refer to source of care within 1 month.	Weight-management counseling if overweight, introduce physical activity and diet management.‡	Initiate therapy based on indications in Table 6 or if compelling indications as above.
Stage 2 hypertension	>99th percentile plus 5 mmHg	Evaluate or refer to source of care within 1 week or immediately if the patient is symptomatic.	Weight-management counseling if overweight, introduce physical activity and diet management.‡	Initiate therapy.§

BP, blood pressure; CKD, chronic kidney disease; DBP, diastolic blood pressure; LVH, left ventricular hypertrophy; SBP, systolic blood pressure

\* For sex, age, and height measured on at least three separate occasions; if systolic and diastolic categories are different, categorize by the higher value.

† This occurs typically at 12 years old for SBP and at 16 years old for DBP.

‡ Parents and children trying to modify the eating plan to the Dietary Approaches to Stop Hypertension (DASH) eating plan could benefit from consultation with a registered or licensed nutritionist to get them started.

§ More than one drug may be required.

TABLE 6

**Indications for Antihypertensive Drug Therapy in Children**

- Symptomatic hypertension
- Secondary hypertension
- Hypertensive target-organ damage
- Diabetes (types 1 and 2)
- Persistent hypertension despite nonpharmacologic measures